

# **Integrated Studies of Oceanographic Processes and Shallow Water Acoustics in the South China Sea: Custom Climatology and Mid-Shelf Field Work**

Glen Gawarkiewicz  
Mail Stop #21  
Woods Hole Oceanographic Institution  
Woods Hole, MA 02543  
phone: (508) 289-2913 fax: (508) 457-2181 email: [gleng@whoi.edu](mailto:gleng@whoi.edu)

Award Number: N00014-04-1-0720  
[http://www.whoi.edu/science/po/people/clinder/projects\\_mabclimatology.html](http://www.whoi.edu/science/po/people/clinder/projects_mabclimatology.html)

## **LONG-TERM GOALS**

Our long-term goal is to understand the dynamics of shelfbreak processes in the South China Sea and the associated forcing from the Kuroshio. We also collaborate with acousticians to determine how these processes affect the propagation of low-frequency sound near the shelfbreak.

## **OBJECTIVES**

The objectives of this past year were to finish a climatology of the northern South China Sea shelfbreak and Taiwan Strait as well as to explore issues relating to internal tides near the shelfbreak as well as instability of the Kuroshio in Luzon Strait.

## **APPROACH**

Our approach has been to use climatological tools developed previously for the Middle Atlantic Bight (Linder et al., 2006) to the shelfbreak in the northern South China Sea, the East China Sea, and in Taiwan Strait. We collapsed the data into a single cross-shelf section in order to establish the cross-shelf structure on the scale of the baroclinic Rossby radius. We have also referenced the geostrophic velocity fields using seasonally averaged data from shipboard ADCP sections collected by the Taiwanese.

## **WORK COMPLETED**

During the past year, we have submitted a manuscript (Linder et al., 2008) on the climatological work. We have been collaborating with J.-H. Tai of National Taiwan University on a stability analysis of the Kuroshio in Luzon Strait. A paper on the impact of the oceanographic variability on transmission loss near the shelfbreak in the South China Sea has appeared (Emerson et al., 2007).

## **RESULTS**

We have computed a four season, two-dimensional climatology for both Taiwan Strait and the northern South China Sea, centered on the shelfbreak (Figure 1). We found a predominantly northeastward flow near the shelfbreak in the South China Sea, consistent with an anti-cyclonic gyre southwest of Taiwan. Maximum velocities over the upper slope are 40 cm/s. The maximum

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2008</b>		2. REPORT TYPE <b>Annual</b>		3. DATES COVERED <b>00-00-2008 to 00-00-2008</b>	
4. TITLE AND SUBTITLE <b>Integrated Studies Of Oceanographic Processes And Shallow Water Acoustics In The South China Sea: Custom Climatology And Mid-Shelf Field Work</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Woods Hole Oceanographic Institution,Mail Stop #21,Woods Hole,MA,02543</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>code 1 only</b>					
14. ABSTRACT <b>Our long-term goal is to understand the dynamics of shelfbreak processes in the South China Sea and the associated forcing from the Kuroshio. We also collaborate with acousticians to determine how these processes affect the propagation of low-frequency sound near the shelfbreak.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

variability occurs just seaward of the shelfbreak in a depth range of 40 to 80 m for temperature, and near the surface at the shelfbreak for salinity. Transports, using mean barotropic velocities from all available shipboard ADCP sections, are at a maximum in summer (1.8 Sv) and at a minimum in spring (0.1 Sv). The seasonally averaged transports for both the shelfbreak in the northern South China Sea as well as Taiwan Strait appears in Figure 2.

In Taiwan Strait, the dominant variability is present on the western side of the strait and is associated with the China Coastal Current. Maximum transport is to the north and is 1.9 Sv in summer (compared to previous estimates of 1.5 Sv) and a minimum of 0.4 Sv in spring, again northward. The mean and standard deviation fields appear in Figures 3 and 4.

J.-H. Tai is proceeding toward his Ph.D. degree at National Taiwan University after having spent a year at Woods Hole Oceanographic Institution working with G. Gawarkiewicz. He has completed a two-layer model instability analysis of the Kuroshio over ridge topography and is presently comparing his results to moored measurements in Luzon Strait. The basic state for the model was chosen based on sections across the Kuroshio taken August, 2007 using the SeaSoar from National Taiwan University.

We are continuing to do analysis of data collected in April, 2005 (Emerson et al., 2007) with attention on the internal tides. We have identified strong internal bores in the data and are presently examining the bore structure as well as boluses of cold water which appear on the shelf.

## **IMPACT/APPLICATIONS**

We have used the climatology to do planning of field data in the East China Sea for the Quantifying, Predicting, and Exploiting Uncertainty Pilot program, including using the climatological sound speed fields for acoustic propagation modeling. We have a technical report which is web-based and has been distributed to several fleet contacts.

## **RELATED PROJECTS**

This work is closely related to the Quantifying, Predicting, and Exploiting Uncertainty DRI. We have used sampling strategies and equipment (low-cost thermistor moorings) directly in the QPE Pilot Program. We have also used the climatological tools to produce climatological fields which have been used to initialize numerical models and for acoustic propagation modeling in the QPE DRI.

## **PUBLICATIONS**


Emerson, C., P. Abbot, C. Gedney, G. Gawarkiewicz, C.-S. Chiu, C.-F. Chen, and R., Wei, "Acoustic Propagation Uncertainty in the Shallow South China Sea", Proceedings, Underwater Acoustic Measurements: Technologies and Results, Papadakis and Bjorno, ed., 25-29 June, 2007.

Linder, C., G. Gawarkiewicz, J.-H. Tai, and T.-Y. Tang, 2008. A seasonal climatology of Taiwan Strait and the northern South China Sea. J. Geophys. Res.-Oceans, submitted.

## FIGURES


QuickTime™ and a  
TIFF (Uncompressed) decompressor  
are needed to see this picture.

***Figure 1- A map denoting the areas used for the climatological fields produced for the northern South China Sea and Taiwan Strait. All data from within the polygon is used and mapped relative to the reference bathymetry (light dashed line). The dark dashed line represents the cross-shelf (NESCO) and cross-strait (TS) section that the data is projected onto.***




**Figure 2- The seasonal variation in transport within the northern South China Sea (dashed line) and Taiwan Strait (solid line).**

# Taiwan Strait: March, April, May



**Figure 3- Mean fields of temperature (upper panel), salinity (middle panel), and density (bottom panel) for spring in Taiwan Strait. The bins used for averaging are denoted at the top of the panels. Note the presence of the China Coastal Current in the salinity field at the left side of the figure (western side of Taiwan Strait).**

Taiwan Strait: March, April, May



**Figure 4- The spatial structure of the variability within Taiwan Strait as expressed in the standard deviation fields. The top panel is temperature, the middle panel is salinity, and the bottom panel is density. The maximum standard deviations appear at the western end of the strait and are associated with the China Coastal Current.**